

IN THE GRAND SCHEME:


FILL-IN: In Photosynthesis plants convert light energy of the sun into chemical bond energy stored in organic molecules. Plants are autotrophic organisms that “feed themselves”, in the sense that they make their own organic molecules from inorganic raw materials. Some bacteria are chemoautotrophs, which means they use energy from oxidizing inorganic substances to produce organic compounds.

REVIEW...

Have each member of your Learning Community, in turn, define the KEY TERMS given below and tell how it may relate to the concepts of energy and metabolism. As a group, help each other answer any questions concerning any portion of the review section that any person does not understand before proceeding further.

- a) **Thylakoid** - a flattened membrane sac inside the chloroplast, used to convert light energy into chemical bond energy. Stacks of thylakoid membranes are called grana.
- b) **Chloroplast** - the fluid (aqueous) portion of the chloroplast surrounding the thylakoid membranes, contains bacterial size ribosomes, DNA, enzymes of CO₂ fixation, and solutes. Also called the **stroma**.
- c) **Calvin Cycle** - a major biochemical pathway of photosynthesis involving the reduction of atmospheric carbon dioxide into carbohydrate, as glucose
- d) **Electromagnetic energy** - the entire spectrum of radiation, produced by the fusion reactions of the sun; expressed in wavelengths, which run from less than a nanometer to more than a kilometer. A portion of the spectrum is referred to as the visible spectrum, which includes the light seen by the human eye, and runs from 340 nm to 720 nm.
- e) **Absorption spectra** - a graphical plot of the amount of light absorbed by a purified form of a molecule vs. the wavelengths of the electromagnetic (visible light)
- f) **PEP Carboxylase** - an enzyme found in C₄ plants, which reduces carbon dioxide by combination of CO₂ with PEP to make malate. The efficiency of PEP carboxylase for CO₂ is much greater than that of RuBP carboxylase, the enzyme of CO₂ reduction in C₃ plants.
- g) **CAM Plant** - a plant that uses Crassulacean acid metabolism (CAM), an adaptation for photosynthesis in arid conditions, first discovered in the plant family Crassulaceae. CO₂ entering open stomata during the night is first converted into organic acids (malate) which releases CO₂ for use in the Calvin cycle during the day, when the stomata are closed, thus these processes are temporally separated.
- h) **Photosystem** - a light harvesting complex located within the thylakoid membranes & consisting of an antennae complex of chlorophyll molecules & a reaction center pigment molecule
- i) **Cyclic electron flow** - a route of electron flow, during the light reactions of photosynthesis, which involves only photosystem I (PSI), produces ATP, but not NADPH, and has the electron returning to its original source pigment - P₇₀₀.
- j) **Photorespiration** - a metabolic pathway that consumes oxygen (like cell respiration) and releases carbon dioxide, generates no ATP, and decreases photosynthetic output of carbohydrate. It occurs most commonly on hot, dry, bright days, when plant stomata are closed and the oxygen concentration in the leaf exceeds that of carbon dioxide in Calvin plants.
- k) **Plastoquinone** - a cytochrome-like component of the photosynthetic electron transfer chain which undergoes a redox reaction by gaining/losing electrons and/or protons.

CONCEPT MAPS: Fill in the following concept map that summarizes this section on Photosynthesis.




- A. light energy
- B. organic molecules
- C. water is split
- D. Calvin cycle
- E. NADPH
- F. ATP
- G. photophosphorylation
- H. regenerate RuBP

MORE FILL-IN THE BLANKS...

Chloroplasts are found primarily in the mesophyll tissues of a leaf, and contain the pigments chlorophyll, carotenes, phycobilins. The gases CO₂ and O₂ enter and leave the leaf through stomata. Veins carry water from the roots to the leaves and distribute sugars made in the leaf to other non-photosynthetic tissues.


LABEL IDENTIFICATIONS:



A. Fill in the blanks in this overview of photosynthesis in a chloroplast.


- a. light
- b. H₂O
- c. light reaction take place in grana
- d. O₂
- e. ATP
- f. NADPH
- g. CO₂
- h. Calvin Cycle in the stroma
- i. CH₂O (sugars)

2. Label the indicated parts in this diagram of the chloroplast.



- a. outer membrane
- b. granum
- c. inner membrane
- d. thylakoid compartment
- e. thylakoid
- f. stroma (chloroplast)

3. Label the absorption spectra and the action spectra in the graph below. What is the difference between these two spectra? On this plot draw an action spectra for rhodopsin, a human visual pigment.



The solid line is the absorption spectra showing absorption maxima at 450nm and 660nm indicating that it is probably a chlorophyll. The dashed line is the action spectra.

An absorption spectra is a plot vs. wavelength of the amount of light absorbed, which an action spectra is a plot vs. physiological activity.

An action spectra for rhodopsin would reveal a maxima of optical nerve activity near 500nm in the green regions of the visible spectrum.

SOME INTERACTIVE THOUGHT QUESTIONS...

1. Why is no oxygen generated by cyclic electron flow?

No electrons are released from PSII (P_{680}) and therefore without an oxidized P_{680} there is no need to split water to re-reduce the P_{680} .

2. During chemiosmosis in chloroplasts the proton gradient across the membranes is as great as 1.0 to 2.0 pH units. Name the side of the membrane with the lowest pH?

Inside the thylakoid compartment (locule) is the site of the lowest pH, greatest H^+ concentrations.

3. What possible explanation is there for photorespiration, a process that can result in the loss of as much as 50% of the carbon dioxide reduced in the Calvin Cycle?


Photorespiration may be an evolutionary relic from the time when there was little oxygen in the atmosphere and the ability of the enzyme Rubisco to distinguish between O_2 and CO_2 was not that critical.

Now in our oxygen rich atmosphere, Photorespiration seems to be a great agricultural liability.

4. Exactly where does the Calvin Cycle take place in C_4 plants? *In the Bundle Sheath cells.*


5. In the figure to the right the isolated chloroplasts were first allowed to equilibrate in an acidic solution at pH 4.0. After the chloroplast's thylakoid compartments reached a pH of 4.0, the chloroplasts were collected and transferred to a basic solution at pH 8.0. This caused the chloroplasts to make ATP, even if placed in the dark. Explain?

The higher concentration of H^+ ions inside the locules after being equilibrated at pH 4.0 means that these H^+ ions will move from the locules, with the existing H^+ ion gradient, back out into the high pH media (low H^+ concentrations) of the stroma, probably through the ATP synthase in these membranes, thereby making ATP.



ELECTRON FLOW IN THYLAKOID MEMBRANES...

Label the diagram below (A. through I.), which depicts the molecules and components of electron flow in photosynthesis.



- hydrogen ions (H^+)
- hydroxyl ions (OH^-)
- ATP
- ATP synthase
- locular space
- NADPH
- NADP Reductase
- thylakoid compartment
- photosystem II

DARK REACTIONS OF PHOTOSYNTHESIS:

A little easier this time. Using the following terms to complete the diagram of the Calvin Cycle given below. A term may be used more than once.

- | | |
|--------------------------------|------------------------|
| 1. $ATP \rightarrow ADP + P_i$ | a. 2 and 6 |
| 2. CO_2 | b. 6 |
| 3. glucose and other sugars | c. 5 |
| 4. $NADPH \rightarrow NADP^+$ | d. & e. 1 and 4 |
| 5. PGA | f. 3 |
| 6. RuBP | g. 1 |

